arXiv:1210.4526v1 [nucl-th] 16 Oct 2012

The influence of strong magnetic fields on proto-quark stars

V. Dexheimer,[1, 2] D. P. Menezes,[1] and M. Strickland[2]
[1]*Depto de Física - CFM - Universidade Federal de Santa Catarina Florianópolis SC - CP. 476 - CEP 88.040 - 900 - Brazil*
[2]*Physics Department, Gettysburg College, Gettysburg, PA 17325 United States*

(Dated: October 17, 2012)

We use the MIT bag model to analyze different stages of magnetized quark star evolution incorporating baryon number conservation and an anisotropic energy momentum tensor. The first stages of the evolution are simulated through the inclusion of trapped neutrinos and fixed entropy per particle, while in the last stage the star is taken to be deleptonized and cold. We find that, although strong magnetic fields allow for more massive quark stars, the evolution of isolated stars needs to be constrained by fixed baryon number, which lowers the star masses. Moreover, magnetic field effects, measured by the difference between the parallel and perpendicular pressures, are more pronounced in the beginning of the star evolution when there is a larger number of charged leptons and up quarks. We also show that having a spatially varying magnetic field allows for larger magnetic fields to be supported since the model employed generates large magnetic fields only at high densities, where the longitudinal matter pressure is large enough to partially compensate for the negative magnetic field longitudinal pressure.

PACS numbers: 26.60.kp, 26.60.+c, 95.30.Tg, 95.85.Sz

I. INTRODUCTION

Neutron stars are compact objects with masses of the order of 1-2 $M_\odot$, radii around 10 km, and a temperature of approximately 10^{11} K at birth, cooling rapidly by emitting neutrinos. In conventional neutron star models, they are comprised of hadronic matter and leptons. Nevertheless, according to the Bodmer-Witten conjecture, the true ground state of matter could be deconfined quarks in environments where densities are very large [1–3]. This hypothesis lead to the possibility that neutron stars could also be strange stars or quark stars, comprised of quarks and leptons [4]. The true nature of these compact stars remains a source of speculation and further constraints are necessary so that models can be ruled out.

Magnetars have extremely high magnetic fields of up to $B = 10^{15}$ G on the surface [5–8]. Nevertheless, all pulsars have relatively strong magnetic fields [9, 10], and for this reason, any complete analysis of pulsar features should include magnetic field effects. Unfortunately, all realistic calculations of magnetic field limits in the center of stars are model dependent to some extent, even when the virial theorem is considered. Some results along this line can be found in Refs. [11–18] and have limits ranging between $B = 10^{17} - 10^{20}$ G. At such high magnetic fields, the thermodynamical as well as the hydrodynamical properties of matter become anisotropic [19–25] and, therefore, need to be carefully handled.

Many pulsars exist in binary systems, however, it is possible to have an isolated pulsar, in which case one needs to take care to properly enforce baryon number conservation. This happens because, differently from pulsars in binary systems which can accrete matter from a companion, isolated stars cannot and, therefore, they conserve baryon number. In this case, as was previously shown in Refs. [26–28], there exist windows of stability which only allow for stars with particular values of baryon number to exist at any given time. The different snapshots of the evolution are simulated though different entropies per particle and trapped neutrinos. As was shown in Ref. [29] for a hadronic model, such an analysis shows that the maximum mass that can possibly be attained by hadronic stars decreases with time.

There have been some real-time calculations of the evolution of proto-neutron stars performed in Refs. [30–33] utilizing Boltzmann transport equations with realistic neutrino cross sections. Although a very interesting preliminary calculation of the neutrino absorption cross section influenced by magnetic field effects was performed in Ref. [34] for neutron stars, to the best of our knowledge, such early time simulations have not been performed including magnetic field effects in quark matter.

There have been previous studies of proto-quark and proto-hybrid stars [35–47]. Others have considered proto-neutron stars subject to a magnetic field [48, 49] and zero temperature quark stars in a magnetic field using the bag model [50, 51]. In this paper we make the first study of the effects of magnetic fields on finite temperature proto-quark stars incorporating the effect of baryon number conservation. This is important when considering isolated proto-quark stars. In addition, we include the pressure anisotropies generated by all relevant degrees of freedom (quarks, leptons, and neutrinos) and the magnetic field.

Using the MIT bag model [52], we determine the effect of strong magnetic fields on three snapshots of quark star evolution. Our task includes a comprehensive investigation of the pressure anisotropy at zero temperature and at fixed entropy per baryon, including finite temperature effects. In most of the previous works on neutron stars subject to strong magnetic fields, the anisotropic pressures were either disregarded [53–57] or only considered at zero temperature [20, 21, 23]. Here we ignore the effects of the quark anomalous magnetic moment (AMM) and focus on the effect of Landau quantization only.

The structure of the paper is as follows. In Sec. II we review the formal setup of the problem and present expressions for the bulk properties of the system. In Sec. III we present our results for the transverse and longitudinal pressures and analyze the effect on the mass-radius relationship and star properties at three different moments in its evolution. In Sec. IV we present our conclusions and an outlook for the future.

II. FORMALISM

The Lagrangian density for the bag model including the magnetic field is given by

$$\mathcal{L} = \left[\bar{\Psi}_q \left(i\gamma^\mu \partial_\mu - e_q \gamma^\mu A_\mu - m_q\right) \Psi_q - \mathcal{B}\right]\Theta_V - \frac{1}{16\pi} F_{\mu\nu}F^{\mu\nu} + \bar{\Psi}_l \left(i\gamma^\mu \partial_\mu - e_l \gamma^\mu A_\mu - m_l\right) \Psi_l, \quad (1)$$

where Ψ_q stands for the 3 light quark fields, $e = 0.08$ is the electron charge (in Gaussian natural units), A^μ is the electromagnetic field and Θ_V is a step function defined as 1 inside the bag and 0 outside. The bag constant value is chosen to be $\mathcal{B} = (154\ \mathrm{MeV})^4$ in accordance with an analysis of stability made for magnetized proto-quark stars [58]. $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field tensor (multiplied by $1/\sqrt{4\pi}$ due to our choice of units). Finally, Ψ_l stands for the electron, muon, and electron and muon neutrino fields.

The magnetic field is taken to be parallel to the z direction. For details of the calculation of the bulk properties of a Fermi gas we refer to Refs. [20, 21, 23, 25, 53–57, 59–62]. The pressure and energy density in the bag model are the pressure and energy density of a non-interacting Fermi gas with the bag constant subtracted and added, respectively. For the charged particles, the matter contribution to the pressure and energy density for finite and zero temperature in the presence of a constant magnetic field, B, in the z direction are

$$P_{m_\parallel} = \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \int \frac{\bar{k}_i^2}{\sqrt{\bar{k}_i^2 + \bar{m}_i^2}} (f_{+i} + f_{-i}) d\bar{k} - \mathcal{B} \quad (2)$$

$$P_{m_\parallel}(T=0) = \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \times \frac{1}{2}\left[\bar{k}_{Fi}\mu_i - \bar{m}_i^2 \ln\left|\frac{\bar{k}_{Fi} + \mu_i}{\bar{m}_i}\right|\right] - \mathcal{B}, \quad (3)$$

$$\epsilon_m = \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \int \sqrt{\bar{k}_i^2 + \bar{m}_i^2} (f_{+i} + f_{-i}) d\bar{k} + \mathcal{B}\,, \quad (4)$$

$$\epsilon_m(T=0) = \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \times \frac{1}{2}\left[\bar{k}_{Fi}\mu_i + \bar{m}_i^2 \ln\left|\frac{\bar{k}_{Fi} + \mu_i}{\bar{m}_i}\right|\right] + \mathcal{B}, \quad (5)$$

where γ_i is the degeneracy of each particle i (6 for quarks, 2 for electrons and muons and 1 for neutrinos) taking into account spin and/or number of colors, Q_{e_i} is the electric charge of each particle (in multiples of the electron charge), B the magnetic field, $\bar{k}_{Fi} \equiv \sqrt{\mu_i^2 - \bar{m}_i^2}$ the Fermi momentum of each particle modified by the magnetic field, $\bar{m}_i = \sqrt{m_i^2 + 2|Q_{e_i}|eB\nu}$ the mass of each particle modified by the magnetic field with m_i being the bare mass of each particle (with $m_{u,d} = 5$ MeV and $m_s = 150$ MeV). The distribution functions for the particles (f_+) and anti-particles (f_-) are $f_{\pm i} = 1/[e^{(\sqrt{\bar{k}_i^2 + \bar{m}_i^2} \mp \mu_i)/T} + 1]$, where T is the temperature.

The chemical potential of each particle μ_i is calculated from the independent chemical potentials (baryon, charge and lepton chemical potentials), each of which is related to a conserved quantity (baryon number, electric charge and lepton fraction)

$$\mu_i = Q_{B_i}\mu_B + Q_{e_i}(\mu_q + \mu_l) + Q_{l_i}\mu_l, \quad (6)$$

where Q_{B_i} is the baryon number (1/3 for quarks and 0 for leptons) and Q_{l_i} the lepton number (0 for quarks and 1 for leptons) of each particle. Note that the lepton chemical potential is non-zero only when the lepton fraction is fixed. If strangeness were conserved, like in the case of the initial stages of a heavy ion collision, a strange chemical potential would have to be introduced.

Due to energy level quantization in directions perpendicular to the magnetic field, the thermodynamic quantities for charged particles are summed over the Landau levels ν, which for zero temperature have a maximum value equal to $\nu_{max} = \lfloor(\mu_i^2 - m_i^2)/(2|Q_{e_i}|eB)\rfloor$. Note that the degeneracy of the zeroth Landau level is always half of the usual value for spin 1/2 particles. For non-charged particles, such as neutrinos, the pressure and energy density expressions take the usual form [52].

For the star calculations, we require that the matter is charge neutral and in chemical equilibrium. The first condition can be written as

$$\sum_i Q_{e_i}\rho_i = 0, \quad (7)$$

where the density of each charged particle for finite and zero temperature cases is

$$\rho_i = \sum_\nu \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \int (f_{+i} - f_{-i}) d\bar{k}, \quad (8)$$

$$\rho_i(T=0) = \sum_\nu \frac{\gamma_i}{2\pi^2} |Q_{e_i}| eB \bar{k}_{Fi}, \quad (9)$$

and the total baryon density is simply $\sum_i Q_{B_i}\rho_i$. The second condition, corresponding to chemical equilibrium, can be written by replacing the baryon number, electric charge, and lepton number for each particle in Eq. (6):

$$\begin{aligned}
\mu_u &= \frac{1}{3}\mu_B + \frac{2}{3}\mu_q + \frac{2}{3}\mu_l, \\
\mu_d &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_q - \frac{1}{3}\mu_l, \\
\mu_s &= \frac{1}{3}\mu_B - \frac{1}{3}\mu_q - \frac{1}{3}\mu_l, \\
\mu_e &= -\mu_q, \\
\mu_\mu &= -\mu_q, \\
\mu_{\nu_e} &= \mu_l, \\
\mu_{\nu_\mu} &= \mu_l. \qquad (10)
\end{aligned}$$

It is easy to see from Eq. (10) that $\mu_d = \mu_s = \mu_u + \mu_e - \mu_l$, $\mu_e = \mu_\mu$, and $\mu_{\nu_e} = \mu_{\nu_\mu}$.

Due to the magnetic field, the perpendicular component of the matter contribution to the pressure is also modified. In fact, one can show that the difference of the parallel and perpendicular components of the pressure is proportional to the magnetization. The magnetization can be calculated by taking the negative of the derivative of the grand potential with respect to the magnetic field, resulting for finite and zero temperature in

$$M = -\frac{\partial \Omega}{\partial B} = \frac{P_{m_\parallel}}{B} - \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} {Q_{e}}_i^2 e^2 B\nu \int \frac{1}{\sqrt{\bar{k}_i^2 + \bar{m}_i^2}} \times (f_{+i} + f_{-i}) d\bar{k}, \qquad (11)$$

$$M(T=0) = \frac{P_{m_\parallel}}{B} - \sum_{i,\nu} \frac{\gamma_i}{2\pi^2} {Q_{e}}_i^2 e^2 B\nu \ln\left|\frac{\bar{k}_{Fi} + \mu_i}{\bar{m}_i}\right|, \qquad (12)$$

so the perpendicular component of the matter contribution to the pressure becomes

$$P_{m_\perp} = P_{m_\parallel} - MB. \qquad (13)$$

Due to the presence of the magnetic field, there are, in addition, contributions from the electromagnetic field tensor (see Eq. (1)) to the pressures and energy density [63, 64]. Due to the breaking of rotational symmetry by the magnetic field, these contributions are again different in the parallel and perpendicular directions. The resulting pressures and energy density are

$$\epsilon = \epsilon_m + \frac{B^2}{8\pi}, \qquad (14)$$

$$P_\perp = P_{m_\perp} + \frac{B^2}{8\pi}, \qquad (15)$$

$$P_\parallel = P_{m_\parallel} - \frac{B^2}{8\pi}, \qquad (16)$$

where the 4π comes from the choice of Gaussian natural units.



FIG. 1. (Color online) Binding energy versus baryon density for charge neutral matter at zero temperature shown with and without (constant) magnetic field effects. For the magnetized case $B = 4.30 \times 10^{18}$ G.

The temperature is not expected to be constant in the interior of compact stars [4]. This can be easily taken into account by computing quantities at fixed entropy per particle. Numerical simulations [30–32, 65] show that this ratio can reach values 1 or 2, depending on the stage of evolution. In this case, the temperature increases naturally from a smaller value on the surface to a higher value in the center of the star. The entropy per particle (baryon) can be calculated through the thermodynamical expression

$$\frac{S}{A} = \frac{s}{\rho_B} = \frac{\epsilon + P - \mu_B \rho_B}{T \rho_B}. \qquad (17)$$

Immediately after the supernova explosion, the electron and muon neutrinos are still trapped, which can be modeled using a fixed lepton fraction defined as

$$Y_l = \frac{\sum_i Q_{l_i} \rho_i}{\rho_B}, \qquad (18)$$

which according to simulations [30] is $Y_l \simeq 0.4$. Here we assume $Y_l = 0.4$ throughout.

III. RESULTS AND DISCUSSIONS

We begin our analysis with the binding energy for matter without leptons. The results at zero temperature are shown in Fig. 1. In this work, as previously stated, we use a bag constant of $\mathcal{B} = (154 \text{ MeV})^4$, which reproduces a minimum in the binding energy ($B/A = 925.9$ MeV) slightly below the one for the iron. We point out that the inclusion of the magnetic field makes matter even more stable, giving a binding energy that is even lower ($B/A = 914.30$ MeV). Such a behavior was already pointed out in Refs. [66, 67].

We now proceed with our analysis of the equation of state of the matter only (the pure magnetic field contributions of $\pm B^2/8\pi$ are not included at first). The parallel and perpendicular pressures of matter are plotted in Fig. 2 as a function of energy density of matter for three different cases:



FIG. 2. (Color online) Different matter contributions to the pressure versus matter contribution to the energy density at different snapshots of the star evolution shown with and without (constant) magnetic field effects. For the magnetized case $B = 4.30 \times 10^{18}$ G.



FIG. 3. (Color online) Parallel (red/orange lines) and perpendicular (respective green/dark green lines) pressures of matter versus magnetic field representing different snapshots of the star evolution.

i) $s/\rho_B = 1$, $Y_l = 0.4$,

ii) $s/\rho_B = 2$, $\mu_{\nu_l} = 0$,

iii) $s/\rho_B = 0$, $\mu_{\nu_l} = 0$.

These correspond to three snapshots of the time evolution of a quark star in its first minutes of life. Such an analysis is important, as it has been shown in calculations/simulations that quark matter can already be formed during supernova explosions [68, 69]. At first, the star is relatively warm (represented by fixed entropy per particle) and has a large number of trapped neutrinos (represented by fixed lepton fraction). As the trapped neutrinos diffuse, they heat up the star [27]. Finally, the star can be considered cold.

Notice in Fig. 2 that (constant) magnetic field effects, seen by the difference between the parallel and perpendicular pressures, are more pronounced in the beginning of the star evolution (i), when there is a larger number of negatively charged leptons and up quarks. Other effects, such as the Haas-van Alphen oscillations [70–73] can only be seen at the last stage of evolution (iii), since such oscillations are smoothed out by thermal effects. These oscillations can be better seen in Fig. 3 where the parallel and perpendicular pressures of matter are plotted as a function of magnetic field for different chemical potentials representing the center of the star ($\mu_B = 1500$ MeV) and an outer region ($\mu_B = 1200$ MeV). Our results for $T = 0$ can be compared with the ones presented in Fig. 1 of Ref. [23]. The models are not the same, but the same physics is obtained for $\mu_B = 1500$ MeV (which corresponds to $\mu_q = 500$ MeV in Ref. [23]). At these high densities, the matter contribution to the perpendicular pressure only deviates significantly from the parallel one when the magnetic fields are as high as 3×10^{19} G.

The magnetic field configuration considered in Figs. 1 and 2 was constant in the z direction with $B = 4.30 \times 10^{18}$ G. One expects a non-constant magnetic field whose magnitude decreases as one goes out from the center towards the edge of the star. To allow for this, we consider a magnetic field with a variable strength $B^*(\mu_B)$ that increases with chemical potential (density). It ranges from a lower value set to $B_{surf} = 10^{15}$ G at small baryon chemical potentials (densities) to a higher value set to $B_c = 4.30 \times 10^{18}$ G at high baryon chemical potentials (densities) following the prescription from Refs. [15, 74–76]

$$B^*(\mu_B) = B_{surf} + B_c \left[1 - e^{b\frac{(\mu_B - 938)^a}{938}}\right], \tag{19}$$

with $a = 2.5$, $b = -4.08 \times 10^{-4}$ and μ_B given in MeV. The variable magnetic field prescription employed only significantly modifies the behavior of the system at high densities and therefore does not change the minimum of the binding energy. As explained in [17], stronger magnetic fields are necessary at larger densities in order to conserve the magnetic flux. For an alternative explanation of how the magnetic field strength increases toward the center of the star see [77].

The effects of considering a variable magnetic field can be seen in Fig 4, where the EOS is shown. They only become visible at high densities. It is important to notice that the magnetic field in the center of the star never reaches B_c, but only $\sim 80 - 90\%$ B_c. This happens because the stars become gravitationally unstable for higher central chemical potentials.

We can ask ourselves if it would make sense to increase the magnetic field further than we have already done. The answer is probably not. Beyond $B_c^{\max} \equiv 4.39 \times 10^{18}$ G the parallel pressure (now including the field contribution) begins to become negative in the center of the star, even using the variable magnetic field prescription. According to Ref. [78] this defines an upper limit for the magnetic field, beyond which the system becomes unstable. Using a constant magnetic field, the upper limit is



FIG. 4. (Color online) Different matter contributions to the pressure versus matter contribution to the energy density at different snapshots of the star evolution shown for variable (chemical potential dependent) magnetic field with $B_c = 4.30 \times 10^{18}$ G.



FIG. 5. (Color online) Parallel (red/orange lines) and perpendicular (respective green/dark green lines) pressures versus magnetic field representing different snapshots of the star evolution. Pure magnetic field contribution $B^2/8\pi$ included

lower, as indicated in Fig. 5. For a chemical potential of $\mu_B = 1500$ MeV the parallel pressure becomes negative for $B = 3.17 \times 10^{18}$ G but for $\mu_B = 1200$ MeV this happens for a lower magnetic field of $B = 1.77 \times 10^{18}$ G. Those numbers were obtained for stage (iii) of the star evolution. In the earlier stages, the parallel pressure becomes negative for slightly higher magnetic fields, $B = 3.37 \times 10^{18}$ G for (i) and (ii) at $\mu_B = 1500$ MeV. As this demonstrates, having a variable magnetic field prescription improves the stability since it produces high magnetic fields only at high densities, where the longitudinal matter pressure is large enough to partially compensate for the negative pure magnetic longitudinal pressure.

Having obtained the EoS for the system including magnetic field effects we can use this as input to the TOV equations [79, 80] as long as $B < B_c^{\max}$. We will use the transverse pressure to solve TOV equation. This should be a good approximation when $P_\perp \simeq P_\parallel$; otherwise, one should also include the breaking of spherical symmetry in the solution of the gravitational metric. We will consider two different central magnetic fields of $B_c = 2 \times 10^{18}$ G and $B_c = 4.3 \times 10^{18}$ G, the latter of which is close to $B_c^{\max}$. The results can be seen in the mass-radius diagram shown in Fig. 6. Without magnetic field effects, the maximum mass decreases with time from (i) to (ii) as the star deleptonizes and becomes more isospin symmetric (due to charge neutrality). Actually, the star has more up than down quarks at (i). From (ii) to (iii) the star's maximum mass decreases again as it becomes cold and loses thermal energy. On the other hand, when the (variable) magnetic field effects are included, the maximum mass increases with time, from (i) to (ii) and from (ii) to (iii). The difference in the possible star masses between different snapshots is also larger and all stars are more massive. The bulk of these effects come from the inclusion of the magnetic field in the pressure and energy density.

The analysis above works only for stars that are in binary systems, and therefore can change baryon number with time. In order to analyze isolated stars, we need to constrain the star baryon number by the one at its first moment of evolution (i). As can be seen in Table I the maximum masses of stars in all cases now decrease with time. This result agrees with similar hadronic calculations [29]. The corresponding radii, on the other hand increase with time. It is interesting to notice that within this analysis, the magnetic field in the center of stars from stages (ii) and (iii) only reaches about $\sim 40-50\%\ B_c$. The data from Table I with fixed baryon number is also represented by symbols in Fig. 6.

It is important to note that we assume that the magnetic field is constant in time in the analysis of different snapshots of the quark star evolution. Such an assumption is reasonable considering that the timescale for the magnetic field decays is of the order of $10^5 - 10^7$ years [81–84]. This should be contrasted with the timescale for the cooling of proto-neutron stars which is on the order of minutes.

IV. CONCLUSIONS AND OUTLOOK

In this paper we have examined the effect of strong magnetic fields on proto-quark stars using the MIT bag model. Three situations were investigated: (i) fixed entropy per baryon $s/\rho_B = 1$ with trapped neutrinos, (ii) $s/\rho_B = 2$ without neutrinos and (iii) a zero temperature system with no neutrinos. The pressure anisotropy inherent to Fermi gases in external magnetic fields was obtained both for fixed magnetic fields and for a scenario where the magnetic field varies with the baryon chemical potential. The conditions for chemical equilibrium be-



FIG. 6. (Color online) Mass-radius diagram for star families representing different snapshots of the star evolution shown without and with (variable) magnetic field. The dots represent the time evolution of the most massive stars considering fixed baryon number. The lower part of the figure is not shown as it exhibits the usual behavior.

	$B=0$			$B\neq0$		
stages	i	ii	iii	i	ii	iii
$M_{max}(M_\odot)$	1.66	1.65	1.65	1.84	1.85	1.88
$M_{max}(M_\odot)$ fixed A	1.66	1.58	1.55	1.84	1.73	1.70
R (km) fixed A	9.19	9.53	9.54	9.14	9.62	9.73

TABLE I. Maximum masses with and without fixing baryon number and corresponding radii for different snapshots of the star's evolution with and without the inclusion of a spatially varying magnetic field. For the case with magnetic field we used $B_c = 2.0 \times 10^{18}$ and $B_c = 4.3 \times 10^{18}$ G.

tween the quarks, leptons, and neutrinos were enforced. In the case of isolated proto-quark stars we incorporated the effect of baryon number conservation during its evolution.

We demonstrated that strong magnetic fields allow for more massive quark stars. However, the evolution of isolated stars needs to be constrained by fixed baryon number, which lowers the star masses. For our final mass vs radius results we studied two different central magnetic fields: $B_c = 2.0 \times 10^{18}$ G and $B_c = 4.3 \times 10^{18}$ G. In both cases the maximum magnetic field generated in the star's interior at stage (i) of the evolution is on the order of $B_{\rm max} \lesssim 0.8\, B_c^{\rm max}$. For the lower value of the magnetic field considered, this places us well below $B_c^{\rm max}$ where the longitudinal pressure of the system becomes negative. In addition, the level of pressure anisotropy in this case is relatively small with $P_\parallel/P_\perp \simeq 0.85$, giving us some confidence in the use of isotropic TOV equations which assume $P_\perp = P_\parallel$. For the larger value of the magnetic field studied, we find that the longitudinal pressure is still positive; however, the level of pressure anisotropy is quite large with $P_\parallel/P_\perp \simeq 0.4$. In this second case, it is questionable whether isotropic TOV is sufficient to draw firm numerical conclusions; however, as we have demonstrated, the qualitative behavior and pattern of evolution observed with this larger magnitude of the magnetic field are similar to those obtained with the smaller magnitude magnetic field.

Additionally, our results show clearly that the MIT bag model for a $\mathcal{B}^{1/4} = 154$ MeV obtained from an investigation of the adequate stability window [58] cannot reproduce the very massive neutron star recently detected [85], not even if very intense magnetic fields are considered. This is perhaps not surprising since most models which are able to describe large mass stars rely on the existence of additional repulsive (vector) interactions. That being said, it is interesting to consider the effect of a magnetic field in the somewhat simple bag model in order to establish a baseline for other models.

Finally, we note that although the results presented imply that quark stars with $B_c = 2\times10^{18}$ G are only mildly anisotropic, larger central magnetic fields cause the pressures to become increasingly anisotropic. Therefore, at such values of the magnetic field one should solve Einstein's equations in an axisymmetric metric which is determined self-consistently from the axisymmetric energy-momentum tensor for the star. Numerical solution for the axisymmetric case is needed and we are currently working towards this goal. Other issues to investigate in the future are, for example, the inclusion of the effect of the anomalous magnetic moment and to further refine the stability windows for the bag model in the presence of a background magnetic field.

ACKNOWLEDGMENTS

V.D. and D.P.M acknowledge support from CNPq/Brazil. M.S. was supported by NSF grant No. PHY-1068765 and the Helmholtz International Center for FAIR LOEWE program.

---

[1] N. Itoh, Prog. Theor. Phys. **44**, 291 (1970).
[2] A. R. Bodmer, Phys. Rev. D **4**, 1601 (1971).
[3] E. Witten, Phys. Rev. D **30**, 272 (1984).
[4] D.P. Menezes, C. Providência and D.B. Melrose - J. Phys. G: Nucl. Part. Phys. **32** (2006) 1081-1095.
[5] B. Paczynski, Acta Astron. **42**, 145 (1992).
[6] C. Thompson and R. C. Duncan, Astrophys. J. **392**, L9 (1992).
[7] C. Thompson and R. C. Duncan, Astrophys. J. **473**, 322 (1996).
[8] A. Melatos, Astrophys. J. Lett. **519**, L77 (1999).

[9] J. H. Taylor, R. N. Manchester, and A. G. Lyne, Astrophys. J. S **88**, 529 (1993).
[10] A. G. Lyne, F. Graham-Smith, Pulsar Astronomy (Cambridge Univ. Press., Cambridge, 2005).
[11] M. Bocquet, S. Bonazzola, E. Gourgoulhon and J. Novak, Astron. Astrophys. **301**, 757 (1995).
[12] C. Y. Cardall, M. Prakash and J. M. Lattimer, Astrophys. J. **554**, 322 (2001).
[13] D. Lai and S. L. Shapiro, Astrophys. J. **383**, 745 (1991).
[14] S. Chakrabarty, D. Bandyopadhyay and S. Pal, Phys. Rev. Lett. **78**, 2898 (1997).
[15] D. Bandyopadhyay, S. Chakrabarty and S. Pal, Phys. Rev. Lett. **79**, 2176 (1997).
[16] A. E. Broderick, M. Prakash and J. M. Lattimer, Phys. Lett. B **531**, 167 (2002).
[17] E. J. Ferrer, V. de la Incera, J. P. Keith, I. Portillo and P. P. Springsteen, Phys. Rev. C **82**, 065802 (2010).
[18] M. Malheiro, S. Ray, H. J. Mosquera Cuesta and J. Dey, Int. J. Mod. Phys. D **16**, 489 (2007).
[19] M. Chaichian, S. S. Masood, C. Montonen, A. Perez Martinez and H. Perez Rojas, Phys. Rev. Lett. **84**, 5261 (2000).
[20] A. Perez Martinez, H. Perez Rojas, H. J. Mosquera Cuesta, M. Boligan and M. G. Orsaria, Int. J. Mod. Phys. D **14**, 1959 (2005).
[21] A. Perez Martinez, H. Perez Rojas and H. Mosquera Cuesta, Int. J. Mod. Phys. D **17**, 2107 (2008).
[22] X. -G. Huang, M. Huang, D. H. Rischke and A. Sedrakian, Phys. Rev. D **81**, 045015 (2010).
[23] L. Paulucci, E. J. Ferrer, V. de la Incera and J. E. Horvath, Phys. Rev. D **83**, 043009 (2011).
[24] A. A. Isayev and J. Yang, Phys. Rev. C **84**, 065802 (2011).
[25] M. Strickland, V. Dexheimer and D. P. Menezes, arXiv:1209.3276 [nucl-th].
[26] T. Takatsuka, Nucl. Phys. A **588**, 365 (1995).
[27] M. Prakash, I. Bombaci, M. Prakash, P. J. Ellis, J. M. Lattimer and R. Knorren, Phys. Rept. **280**, 1 (1997).
[28] D. Gondek-Rosinska, P. Haensel and J. L. Zdunik, astro-ph/0012543.
[29] V. Dexheimer and S. Schramm, Astrophys. J. **683**, 943 (2008).
[30] A. Burrows and J. M. Lattimer, Astrophys. J. **307**, 178 (1986).
[31] A. Pons, A. W. Steiner, M. Prakash, and J. M. Lattimer, Phys. Rev. Lett. **86**, 5223 (2001).
[32] J. A. Pons, S. Reddy, M. Prakash, J. M. Lattimer, and J. A. Miralles, Astrophys. J. 513, 780 (1999).
[33] M. D. Alloy and D. P. Menezes, Phys. Rev. **C 83**, 035803 (2011).
[34] T. Maruyama, N. Yasutake, M-K. Cheoun, J. Hidaka, T. Kajino, G. J. Mathews and C-Y. Ryu, arXiv:1201.3455v1 [astro-ph.HE].
[35] A. W. Steiner, M. Prakash and J. M. Lattimer, Phys. Lett. B **509**, 10 (2001).
[36] J. A. Pons, A. W. Steiner, M. Prakash and J. M. Lattimer, Phys. Rev. Lett. **86**, 5223 (2001).
[37] V. K. Gupta, A. Gupta, S. Singh and J. D. Anand, Int. J. Mod. Phys. D **12**, 583 (2003).
[38] J. Berdermann, D. Blaschke, H. Grigorian and D. N. Voskresensky, astro-ph/0512655.
[39] J. -y. Shen, Y. Zhang, B. Wang and R. -K. Su, Int. J. Mod. Phys. A **20**, 7547 (2005).
[40] O. E. Nicotra, M. Baldo, G. F. Burgio and H. -J. Schulze, Phys. Rev. D **74**, 123001 (2006).
[41] G. F. Burgio and S. Plumari, Phys. Rev. D **77**, 085022 (2008).
[42] N. Yasutake and K. Kashiwa, Phys. Rev. D **79**, 043012 (2009).
[43] N. Yasutake, T. Maruyama and T. Tatsumi, J. Phys. Conf. Ser. **312**, 042027 (2011).
[44] I. Bombaci, D. Logoteta, C. Providencia and I. Vidana, arXiv:1102.1665 [astro-ph.SR].
[45] G. Lugones and A. G. Grunfeld, Phys. Rev. D **84**, 085003 (2011).
[46] G. -y. Shao, Phys. Lett. B **704**, 343 (2011).
[47] H. Chen, M. Baldo, G. F. Burgio and H. -J. Schulze, Phys. Rev. D **86**, 045006 (2012).
[48] S. Pal, D. Bandyopadhyay and S. Chakrabarty, astro-ph/9806356.
[49] I. Bednarek, A. Brzezina, R. Manka and M. Zastawny-Kubica, Nucl. Phys. A **716**, 245 (2003).
[50] A. A. Isayev and J. Yang, arXiv:1210.3322 [hep-ph].
[51] S. Chakrabarty, Phys. Rev. D **54**, 1306 (1996).
[52] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. D **9**, 3471 (1974).
[53] D.P. Menezes, M. Benghi Pinto, S.S. Avancini, and C. Providência, Phys. Rev. **C 80**, 065805 (2009).
[54] D.P. Menezes, M. Benghi Pinto, S.S. Avancini, A. Pérez Martinez and C. Providência, Phys. Rev.**C 79**, 035807 (2009).
[55] L.L. Lopes and D.P.Menezes, Braz. J. Phys. (2012), in press, arXiv: 1202.5016 [nuch-th].
[56] A. Rabhi, C. Providencia and J. Da Providencia, J. Phys. G: Nucl. Part. Phys. **35**, 125201 (2008).
[57] A. Rabhi et al, J. Phys. G: Nucl. Part. Phys. **36**, 115204 (2009).
[58] J.R. Torres, D.F. T. Agudelo, V. Dexheimer and D.P. Menezes, in preparation.
[59] H. -Y. Chiu, V. Canuto and L. Fassio-Canuto, Intense Magnetic Fields," Phys. Rev. **176**, 1438 (1968).
[60] V. Canuto and H. Y. Chiu, Phys. Rev. **173**, 1210 (1968).
[61] V. Canuto and H. Y. Chiu, Phys. Rev. **173**, 1220 (1968).
[62] V. Canuto and H. Y. Chiu, Phys. Rev. **173**, 1229 (1968).
[63] J. I. Kapusta and C. Gale, *Finite Temperature Field Theory - Principles and Appications*, Cambridge University Press, 2006.
[64] R. D. Blandford, L. Hernquist, J. Phys. C **15**, 6233 (1982).
[65] J. Stein and J. C. Wheeler, Astrophys. J. **643**, 1190 (2006).
[66] R. G. Felipe and A. P. Martinez, J. Phys. G **36**, 075202 (2009).
[67] X. -J. Wen, S. -Z. Su, D. -H. Yang and G. -X. Peng, Phys. Rev. D **86**, 034006 (2012).
[68] I. Sagert, T. Fischer, M. Hempel, G. Pagliara, J. Schaffner-Bielich, A. Mezzacappa, F. -K. Thielemann and M. Liebendorfer, Phys. Rev. Lett. **102**, 081101 (2009).
[69] T. Fischer, D. Blaschke, M. Hempel, T. Klahn, R. Lastowiecki, M. Liebendorfer, G. Martinez-Pinedo and G. Pagliara *et al.*, Phys. Atom. Nucl. **75**, 613 (2012).
[70] W. J. de Haas and P. M. van Alphen, Leiden Commun. A **212**, 215 (1930).
[71] W. J. de Haas and P. M. van Alphen, Proc. R. Acad. Sci. Amsterdam **33**, 1106 (1930).

[72] D. Ebert, K. G. Klimenko, M. A. Vdovichenko and A. S. Vshivtsev, Phys. Rev. D **61**, 025005 (2000).

[73] D. Ebert and K. G. Klimenko, Nucl. Phys. A **728**, 203 (2003).

[74] G. -J. Mao, A. Iwamoto and Z. -X. Li, Chin. J. Astron. Astrophys. **3**, 359 (2003).

[75] A. Rabhi, H. Pais, P. K. Panda and C. Providencia, J. Phys. G G **36**, 115204 (2009).

[76] V. Dexheimer, R. Negreiros and S. Schramm, arXiv:1108.4479 [astro-ph.HE].

[77] M. Eto, K. Hashimoto and T. Hatsuda, arXiv:1209.4814 [hep-ph].

[78] M. Sinha and B. Mukhopadhyay, arXiv:1005.4995 [astro-ph.HE].

[79] R. C. Tolman, Phys. Rev. **55**, 364 (1939).

[80] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. **55**, 374 (1939).

[81] A. K. Harding and D. Lai, Rept. Prog. Phys. **69**, 2631 (2006).

[82] D. N. Aguilera, J. A. Pons and J. A. Miralles, Astron. Astrophys. **486**, 255 (2008).

[83] J. A. Pons and U. Geppert, Astron. Astrophys. **470**, 303 (2007).

[84] Y. Xie and S. -N. Zhang, arXiv:1110.3869 [astro-ph.HE].

[85] P. B. Demorest et al, Nature **467**, 1081 (2010).